

REPTILIA: SQUAMATA: SAURIA: ANGUIDAE **GERRHONOTUS MULTICARINATUS****Catalogue of American Amphibians and Reptiles.**LAIS, P. MIKE. 1976. *Gerrhonotus multicarinatus*.***Gerrhonotus multicarinatus* (Blainville)
Southern alligator lizard**

Cordylus (*Gerrhonotus*) *multi-carinatus* Blainville, 1835:37. Type-locality, "Californie," restricted to Monterey, [Monterey County] by Fitch (1934b:173). Holotype, adult, Muséum National d'Histoire Naturelle 2002, collected by P.E. Botta (holotype examined by author).

Elgaria multicarinata: Gray, 1838:391.

Gerrhonotus caeruleus: Boulenger, 1885:273 (part).

Gerrhonotus scincicauda: Stejneger, 1893:195.

Gerrhonotus multi-carinatus: Fitch, 1934b:172. Revalidation of *multicarinatus*.

• **CONTENT.** Five subspecies are recognized: *multicarinatus*, *webbii*, *scincicauda*, *ignavus*, and *nanus*.

• **DEFINITION.** A large species of *Gerrhonotus* (maximum snout-vent length 175 mm., tail length over twice snout-vent length), with dorsal scales in 14 longitudinal rows and 40 to 66 transverse rows, and undivided interoccipital. Dorsal scales of both body and tail are heavily keeled. Ventral scales are in 12 longitudinal rows. Dorsal ground color varies from light brown to olive gray or dull yellow, with the sides darker. The body and tail are marked with irregular transverse bands, black on the sides and lighter dorsally, often including scales with white posterior edges. The head is nearly uniform brown with little marking, except for a dark temporal band in some individuals. The venter varies from pale gray to dull yellow, often with longitudinal dark lines aligned in the centers of the scale rows.

• **DESCRIPTIONS.** The original description (Blainville, 1835) is inadequate. Good general descriptions are in Van Denburgh (1922), Fitch (1938), Smith (1946), and Stebbins (1954, 1966). Other descriptions, mainly of taxonomic value, are in Skilton (1849), Baird and Girard (1852, 1853), Baird (1858), Girard (1858), Bocourt (1878), Van Denburgh (1897, 1898), and Cope (1900). The karyotype ($2n = 48$) was described by Bury et al. (1969).


• **ILLUSTRATIONS.** Black and white photographs of adults are in Van Denburgh (1922), Fitch (1935), Gordon (1939), Smith (1946), Pope (1955), Blackstone (1957, mating pair), Banta (1963, bifid tail), and Dixon (1967). Stebbins (1954) has black and white drawings of adult and juvenile, and Stebbins (1966) a color drawing of an adult. Color photographs are in Schmidt and Inger (1957) and Leviton (1972). Line drawings of scutellation are provided by Girard (1858), Bocourt (1878), and Cope (1900).

• **DISTRIBUTION.** *Gerrhonotus multicarinatus* occurs chiefly west of the Cascade-Sierran crest from northern Oregon to northern Baja California. East of the Cascades the range extends along the Columbia River into south-central Washington and north-central Oregon. It occurs on the southern California islands of Anacapa, Santa Cruz, Santa Catalina, Santa Rosa, San Miguel, and San Nicholas; and in Baja California on the islands of Los Coronados and San Martin. In Inyo County, California isolated populations occur east of the Sierra Nevada at Walker Creek near Olancho and southwest of Independence; and in the Mojave Desert along the Mojave River. Throughout its range this species generally occurs in the oak woodlands of low hills and valleys. Occasionally it may be found in mountains, associated with yellow pine, black oak, big-leaf maple, and madrone, but it does not frequent dense coniferous forest. In central and southern California it sometimes occurs near the beach, and may be found under driftwood. Photographs of *G. multicarinatus* habitat are in Fitch (1935).

• **FOSSIL RECORD.** Brattstrom (1955) assigned a frontal bone from Pleistocene deposits of Carpinteria, California to *G. multicarinatus*. Robinson and Van Devender (1973) described Miocene anguids from Wyoming and Nevada as similar to both *G. liocephalus* and *G. multicarinatus*.

• **PERTINENT LITERATURE.** The best taxonomic review is by Fitch (1938); others include Tihen (1949), Stebbins (1958), and Waddick (1963). Natural history notes are in Grinnell et al. (1930), Gander (1931), Fitch (1935), Christenson (1947), Knowlton (1949), Atsatt (1952), Cohen and Wood (1953), Cunningham (1955, 1956), Pope (1955), Montanucci (1968), and Brodie et al. (1969). Studies of reproduction include Burrage (1964, 1965), Greer (1967), Fitch (1970), and Goldberg (1972). Descriptions of eggs and young are in

Shaw (1943, 1952), and Stebbins (1954). Stebbins (1944) reported a specimen killed by a millipede, and Brodie (1968) noted death from *Taricha* skin toxin. Parasitism by ticks was recorded by Mohr et al. (1964), and by a cestode by Telford (1965). Temperature data are in Brattstrom (1965), and notes on physiological responses to temperature in Dawson and Templeton (1966), Cunningham (1966), Regal (1966), and Dawson (1967). Hoffman (1973) discussed locomotor activity and metabolic scope, and Schultz and Norberg (1970) the effect of visual loss on activity. Endocrine system morphology has been studied by Retzlaff (1949, adrenal), Lynn and Colorigh (1967, thyroid), H. Saint Girons (1967, hypophysis), Gabe et al. (1964, adrenal), and Gabe (1970, adrenal). Brain morphology was described by Senn and Northcutt (1973),



MAP. Hollow symbols represent known localities, solid symbols mark type-localities. The star indicates a fossil record.

and brain serotonin content by Quay and Wilhoft (1964). Studies of ear morphology and auditory sensitivity are by Crowley (1964), Wever (1965), Miller (1966), Peterson (1966), Miller et al. (1967), Campbell (1969), and Miller (1973). Stebbins (1948) discussed nasal structure and function. Blood cell morphology and blood chemistry are treated by Dawson and Poulson (1962), Pough (1969), Saint Girons and Saint Girons (1969), Dessauer (1970), and M. Saint Girons (1970). Studies of dentition are in Camp (1923), Edmund (1969), and Soule (1970), and of osteology in Cope (1892), Camp (1923), Etheridge (1967), Criley (1968), Enlow (1969), and Meszoe (1970). Jamison (1964) studied tail regeneration, and McDowell and Bogert (1954) figured the tongue. Peabody and Savage (1958) placed *G. multicarinatus* in a phylogenetic line that moved northward through the Pacific Coast Range Corridor. Distributional notes include Yarrow and Henshaw (1878), Cope (1883), Stejneger (1893), Van Denburgh (1905, 1922), Van Denburgh and Slevin (1914, 1921), Grinnell and Camp (1917), Stephens (1921), Grinnell et al. (1930), Fitch (1936, 1938), Smith and Taylor (1950), Dunlap (1959), Slater (1963), Banta and Morafka (1966), Storm (1966), Bogert and Porter (1967), Bury (1970), and Bostic (1971).

• **ETYMOLOGY.** The name *multicarinatus* (Latin, many keels) refers to the keeled dorsal scales; *webbii* honors Dr. T. H. Webb, collector of the type-specimen; *scincicauda* (Latin, lizard tail) refers to the long tail; the significance of *ignavus* (Latin, lazy or idle) as used by Van Denburgh (1905) is unclear; *Nanus* (Greek, dwarf) refers to the size of this form.

1. *Gerrhonotus multicarinatus multicarinatus* (Blainville) California alligator lizard

Cordylus (*Gerrhonotus*) *multi-carinatus* Blainville, 1835:57. See species account.

Gerrhonotus wiegmanni Gray, 1845:54. Type-locality, "Mexico?," in error, revised to the north side of San Francisco Bay or the lower Sacramento Valley, California, by Fitch (1938:395). Holotype, adult, British Museum (Natural History) 1946.8.7.87, collected by George Barclay (holotype examined by author).

Gerrhonotus scincicauda scincicauda: Grinnell and Camp, 1917:166 (part). First use of trinomial.

Gerrhonotus multi-carinatus multi-carinatus: Fitch, 1934b:173.

Elgaria multicarinata multicarinata: Tihen, 1949:595.

• **DIAGNOSIS.** Maximum snout-vent length about 137mm.; dorsal surface of head often dark-mottled; median row of red blotches on back; body bands dark brown, one scale row wide; one to three rows of scales on upper arm weakly keeled; eight rows of dorsal caudals distinctly keeled near base of tail, sometimes one or two rows of lateral caudals keeled on each side; upper two rows of temporals convex and bluntly keeled or smooth.

2. *Gerrhonotus multicarinatus webbii* Baird San Diego alligator lizard

Gerrhonotus webbii Baird, 1858:255. Type-locality, "near San Diego," San Diego County, California. Holotype, U. S. Natl. Mus. 3078, collected by Dr. T. H. Webb in April or May, 1852? (holotype examined by author).

Gerrhonotus scincicauda webbii: Grinnell and Camp, 1917:168. First use of trinomial.

Gerrhonotus multi-carinatus webbii: Fitch, 1934b:173. First use of combination.

Elgaria multicarinata webbii: Tihen, 1949:596.

• **DIAGNOSIS.** Maximum snout-vent length about 175 mm.; head markings and temporal band absent; bands on back black or dark brown, one to two scale rows wide; three to four rows of keeled scales on upper arm; eight rows of dorsal caudals distinctly keeled near base of tail, several rows of lateral caudals on each side weakly keeled (mean 12.8 rows); upper two rows of temporals heavily keeled.

• **REMARKS.** Plate XXIV, figs. 1-10 in Baird (1859) does not represent the type-specimen of *G. m. webbii*, as might be inferred. It appears to be a juvenile *G. kingi* or *G. panamintinus*, either of which could have been collected on the survey.

3. *Gerrhonotus multicarinatus scincicauda* (Skilton) Oregon alligator lizard

Tropidolepis scincicauda Skilton, 1849:202. Type-locality, "The Dalles of the Columbia," Wasco County, Oregon. Present location of holotype unknown, collected by the Rev. Mr. Gary.

Elgaria grandis Baird and Girard, 1852:176. Type-locality, "Oregon." Holotype, U. S. Natl. Mus. 9057, collected by the U. S. Exploring Expedition under the command of Capt. Charles Wilkes in 1841 (holotype examined by author).

Elgaria scincicauda: Baird and Girard, 1853:348.

Gerrhonotus grandis: Cope, 1875:46.

Gerrhonotus scincicaudus: Cope, 1875:47.

Gerrhonotus scincicauda scincicauda: Grinnell and Camp, 1917:166. First use of trinomial.

Gerrhonotus multi-carinatus scincicauda: Fitch, 1934b:173.

First use of combination.

Elgaria multicarinata scincicauda: Tihen, 1949:595.

• **DIAGNOSIS.** Maximum snout-vent length about 141 mm.; no dark markings on head; bands on back black, one scale row wide; scales on upper arm smooth; eight rows of dorsal caudals keeled, lateral caudals smooth; upper two rows of temporals slightly keeled or smooth.

4. *Gerrhonotus multicarinatus ignavus* Van Denburgh San Martin alligator lizard

Gerrhonotus scincicauda ignavus Van Denburgh, 1905:19. Type-locality, "San Martin Island, Lower California, Mexico." Holotype, California Acad. Sci. 4699, collected by R. H. Beck on 3 May 1903 (holotype examined by author).

Gerrhonotus scincicauda webbii: Grinnell and Camp, 1917:168 (part).

Gerrhonotus multi-carinatus webbii: Fitch, 1938:395 (part).

Gerrhonotus multicarinatus ignavus: Murray, 1955:41. First use of combination.

• **DIAGNOSIS.** Maximum snout-vent length about 124 mm.; brown markings on top of head and on labials, temporal band present; body bands black laterally, black or brown middorsally with a tendency for suffusion of brown pigment over light scales between bands; four to five rows of scales keeled on upper arm; eight rows of dorsal caudals keeled near base of tail, often several rows of lateral caudals keeled on each side (mean 13.4 rows); three to four upper rows of temporals keeled.

5. *Gerrhonotus multicarinatus nanus* Fitch Coronados alligator lizard

Gerrhonotus scincicauda ignavus?: Van Denburgh, 1905:18.

Gerrhonotus scincicauda ignavus: Van Denburgh and Slevin, 1914:140 (part).

Gerrhonotus scincicauda webbii: Van Denburgh, 1922:485 (part).

Gerrhonotus scincicauda nanus Fitch, 1934a:7. Type-locality, "South Island, Los Coronados Islands, Lower California," Mexico. Holotype, adult, Univ. California Mus. Vert. Zool. 5402, collected by A. B. Howell on 1 July 1913 (holotype examined by author).

Gerrhonotus multi-carinatus nanus: Fitch, 1938:397. First use of combination.

Elgaria multicarinata nana: Tihen, 1949:596.

• **DIAGNOSIS.** Maximum snout-vent length about 114 mm.; head marked with dark brown, broad temporal band present; body bands similar to those in *ignavus*; one to three rows of scales on upper arm weakly keeled; eight rows of dorsal caudals strongly keeled near base of tail, two or three rows of lateral caudals on each side usually keeled; upper two rows of temporals sharply keeled, lower ones weakly keeled or smooth.

LITERATURE CITED

- Atsatt, S.R. 1952. Observations on the early life history of the lizards *Sceloporus graciosus vandenburgianus* and *Gerrhonotus multicarinatus webbii*. *Copeia* 1952(4):276.
- Baird, S. F. 1858. Description of new genera and species of North American lizards in the museum of the Smithsonian Institution. *Proc. Acad. Natur. Sci. Philadelphia* 10:253-256.
- 1859. Reptiles of the boundary. In Vol. 2, Pt. 2, United States and Mexican Boundary Survey, U. S. 34th Congress, 1st Session, Exec. Doc. 108:1-35, 41 pls.
- , and C. Girard. 1852. Descriptions of new species of reptiles, collected by the U. S. Exploring Expedition under the

- command of Capt. Charles Wilkes, U.S.N. Proc. Acad. Natur. Sci. Philadelphia 6(5):174-177.
- 1853. Reptiles. In Exploration and survey of the Valley of the Great Salt Lake of Utah by H. Stansbury, U. S. 32nd Congress, Special Session, Exec. Doc. 3, Appendix C:336-365, pls. 1-7.
- Banta, B.H. 1963. An anomalous California red-backed alligator lizard. *Turtos News* 41(8):207.
- , and D. Morafka. 1966. An annotated check list of the recent amphibians and reptiles inhabiting the city and county of San Francisco, California. *Wasmann J. Biol.* 24(2):223-238.
- Blackstone, R. E. 1957. Courtship of the red-backed alligator lizard. *Nature Magazine*, New York 50:312.
- Blainville, M. H. D. de. 1835. Description de quelques espèces de la Californie. *Nouv. Ann. Mus. Nat. Hist. Natur.*, Paris 4:1-64, pls. 24-27.
- Bocourt, M.-F. 1878. Etudes sur les reptiles et les batraciens. In Duméril, Bocourt and Mocquard, *Recherches zoologiques pour servir à l'histoire de la faune de l'Amérique Central et du Mexique*. Miss. Sci. Mexique et Amer. Cent., Imprimerie Nationale, Paris. Part 3, 5:281-360, pls. 20D-G, 21 A-C.
- Bogert, C. M., and A. P. Porter. 1967. A new species of *Abronia* (Sauria, Anguillidae) from the Sierra Madre del Sur of Oaxaca, Mexico. *Amer. Mus. Novitates* (2279):1-21.
- Bostic, D. L. 1971. Herpetofauna of the Pacific Coast of north central Baja California, Mexico, with a description of a new subspecies of *Phyllodactylus xanti*. *Trans. San Diego Soc. Natur. Hist.* 16(10):237-264.
- Boulenger, G. A. 1885. Catalogue of the lizards in the British Museum (Natural History). Second edition. Taylor and Francis, London. Vol. 2: xiii + 497 p.
- Brattstrom, B. H. 1955. Small herpetofauna from the Pleistocene of Carpinteria, California. *Copeia* 1955(2):138-139.
- 1965. Body temperatures of reptiles. *Amer. Midland Natur.* 73(2):376-422.
- Brodie, E. D., Jr. 1968. Investigations on the skin toxin of the adult rough-skinned newt, *Taricha granulosa*. *Copeia* 1968(2):307-313.
- , R. A. Nussbaum, and R. M. Storm. 1969. An egg-laying aggregation of five species of Oregon reptiles. *Herpetologica* 16(3):223-227.
- Burrage, B. R. 1964. The eggs and young of *Gerrhonotus multicarinatus nanus* Fitch. *Herpetologica* 20(2):133.
- 1965. Notes on the eggs and young of the lizards *Gerrhonotus multicarinatus webbi* and *G. m. nanus*. *Copeia* 1965(4):512.
- Bury, R. B. 1970. A biogeographic analysis of the herpetofauna of Trinity County, California. *J. Herpetol.* 4(3-4):165-178.
- , G. C. Gorman, and J. F. Lynch. 1969. Karyotypic data for five species of anguillid lizards. *Experientia* 25(3):314-316.
- Camp, C. L. 1923. Classification of the lizards. *Bull. Amer. Mus. Natur. Hist.* 48(11):289-481.
- Campbell, H. W. 1969. The effects of temperature on the auditory sensitivity of lizards. *Physiol. Zool.* 42(2):183-210.
- Christenson, A. K. 1947. Feeding habits of *Gerrhonotus*. *Herpetologica* 4(1):2.
- Cohen, N. H., and S. F. Wood. 1953. Vertebrate census of an earth, stone, concrete check dam at the San Joaquin Experimental Range. *Bull. Southern California Acad. Sci.* 52(1):35-37.
- Cope, E. D. 1875. Check-list of North American Batrachia and Reptilia; with a systematic list of the higher groups, and an essay on geographical distribution. Based on the specimens contained in the U. S. National Museum. U. S. Nat. Mus. Bull. (1):1-104.
- 1883. Notes on the geographical distribution of Batrachia and Reptilia in western North America. *Proc. Acad. Natur. Sci. Philadelphia* 35(1):10-35.
- 1892. The osteology of the Lacertilia. *Proc. Amer. Philos. Soc.* 30(137):184-221.
- 1900. The crocodilians, lizards and snakes of North America. *Ann. Rept. U. S. Nat. Mus.* for 1898:153-1294.
- Criley, B. B. 1968. The cranial osteology of gerrhonotiform lizards. *Amer. Midland Natur.* 80(1):199-219.
- Crowley, D. E. 1964. Auditory responses in the alligator lizard. *J. Audit. Res.* 4:135-143.
- Cunningham, J. D. 1955. Arboreal habits of certain reptiles and amphibians in southern California. *Herpetologica* 11(3):217-220.
- 1956. Food habits of the San Diego alligator lizard. *Ibid.* 12(3):225-230.
- 1966. Thermal relations of the alligator lizard *Gerrhonotus multicarinatus webbi*. *Ibid.* 22(1):1-7.
- Dawson, W. R. 1967. Interspecific variation in physiological responses of lizards to temperature. P. 230-257. In W. W. Milstead (ed.), *Lizard ecology: A symposium*. University of Missouri Press, Columbia.
- , and T. L. Poulson. 1962. Oxygen capacity of lizard bloods. *Amer. Midland Natur.* 68(1):154-164.
- , and J. R. Templeton. 1966. Physiological responses to temperature in the alligator lizard, *Gerrhonotus multicarinatus*. *Ecology* 47(5):757-765.
- Dessauer, H. C. 1970. Blood chemistry of reptiles: Physiological and voluntary aspects. P. 1-72. In C. Gans and T. S. Parsons (eds.), *Biology of the Reptilia, Morphology C*, Vol. 3. Academic Press, New York.
- Dixon, J. R. 1967. Amphibians and reptiles of Los Angeles County, California. Los Angeles Co. Mus. Natur. Hist., Sci. Ser. (23):1-63.
- Dunlap, D. G. 1959. Notes on the amphibians and reptiles of Deschutes County, Oregon. *Herpetologica* 15(4):173-177.
- Edmund, A. G. 1969. Dentition. P. 117-200. In C. Gans and T. S. Parsons (eds.), *Biology of the Reptilia, Morphology A*, Vol. 1. Academic Press, New York.
- Estes, R. 1964. Fossil vertebrates from the Late Cretaceous Lance Formation, eastern Wyoming. *Univ. California Publ. Geol. Sci.* 49:1-180.
- Etheridge, R. 1967. Lizard caudal vertebrae. *Copeia* 1967(4):699-721.
- Fitch, H. S. 1934a. New alligator lizards from the Pacific Coast. *Copeia* 1934(1):6-7.
- 1934b. A shift of specific names in the genus *Gerrhonotus*. *Copeia* 1934(4):172-173.
- 1935. Natural History of the alligator lizards. *Trans. Acad. Sci. St. Louis* 29:1-38.
- 1936. Amphibians and reptiles of the Rogue River Basin, Oregon. *Amer. Midland Natur.* 17(3):634-652.
- 1938. A systematic account of the alligator lizards (*Gerrhonotus*) in the western United States and Lower California. *Ibid.* 20(2):381-424.
- 1970. Reproductive cycles of lizards and snakes. *Univ. Kansas Mus. Natur. Hist., Misc. Publ.* (52):247 p.
- Gabe, M. 1970. The adrenal. P. 263-318. In C. Gans and T. S. Parsons (eds.), *Biology of the Reptilia, Morphology C*, Vol. 3. Academic Press, New York.
- , M. Martoja, and H. Saint Girons. 1964. État actuel des connaissances sur la glande surrénale des reptiles. *Année Biol.* 3(7-8):303-376.
- Gander, F. 1931. Observations on an alligator lizard. *Copeia* 1931(1):14-15.
- Girard, C. 1858. Herpetology. In *United States Exploring Expedition during the years 1838, 1839, 1840, 1841, 1842, under the command of Charles Wilkes, U. S. N.* J. B. Lippincott, Philadelphia. Vol. 20:xvii + 496 p.
- Goldberg, S. R. 1972. Reproduction in the southern alligator lizard *Gerrhonotus multicarinatus*. *Herpetologica* 28(3):267-273.
- Gordon, K. 1939. The Amphibia and Reptilia of Oregon. *Oregon State Monogr.* (1):1-82.
- Gray, J. E. 1838. Catalogue of the slender-tongued saurians, with descriptions of many new genera and species. Part 2. *Ann. Mag. Natur. Hist.* (1)1(5):388-394.
- 1845. Catalogue of the specimens of lizards in the collection of the British Museum. London. xxvii + 289 p.
- Greer, A. E. 1967. Notes on the mode of reproduction in anguillid lizards. *Herpetologica* 23(2):94-99.
- Grinnell, J., and C. L. Camp. 1917. A distributional list of the amphibians and reptiles of California. *Univ. California Publ. Zool.* 17(10):127-208.
- , J. Dixon, and J. M. Linsdale. 1930. Vertebrate natural history of a section of northern California through the Lassen Peak region. *Univ. California Publ. Zool.* 35:v + 594 p.
- Hoffmann, M. 1973. Locomotor activity, body temperature, and metabolic scope in the southern alligator lizard *Gerrhonotus multicarinatus webbi* (Anguillidae). M. A. Thesis, California State University, Fullerton. 41 p.
- Jamison, J. P. 1964. Regeneration subsequent to intervertebral amputation in lizards. *Herpetologica* 20(3):145-149.
- Knowlton, G. F. 1949. Insect food of the red-backed alligator lizard. *Herpetologica* 5(2):46-47.
- Leviton, A. E. 1972. Reptiles and amphibians of North America. Doubleday and Co., New York. 250 p.
- Lynn, W. G., and M. L. Colorigh. 1967. Thyroid morphology in lizards of the families Anguillidae, Chamaleontidae, and Cordylidae. *Amer. Midland Natur.* 77(1):247-250.
- McDowell, S. B., Jr., and C. M. Bogert. 1954. The systematic position of *Lanthanotus* and the affinities of the anguillid lizards. *Bull. Amer. Mus. Natur. Hist.* 105(1):1-142.

- Mesozoely, C. A. M. 1970. North American fossil anguillid lizards. *Bull. Mus. Comp. Zool.* 139(2):87-147.
- Miller, M. R. 1966. The cochlear duct of lizards. *Proc. California Acad. Sci.* 4th Ser. 33(11):255-359.
- 1973. Scanning electron microscopic studies of some lizard ear basilar papillae. *Amer. J. Anat.* 138(3):301-330.
- , M. Kasahara, and M. Mulroy. 1967. Observations on the structure of the cochlear duct limbus of reptiles. *Proc. California Acad. Sci.* 4th Ser. 35(3):37-52.
- Mohr, C. O., D. E. Beck, and E. P. Brinton. 1964. Observations on host-parasite relationships and seasonal history of ticks in San Mateo County, California. *Great Basin Natur.* 24(1):1-6.
- Montanucci, R. R. 1968. Notes on the distribution and ecology of some lizards in the San Joaquin Valley, California. *Herpetologica* 24(4):316-320.
- Murray, K. F. 1955. Herpetological collections from Baja California. *Herpetologica* 11(1):33-48.
- Peabody, F. E., and J. M. Savage. 1958. Evolution of a Coast Range Corridor in California and its effect on the origin and dispersal of living amphibians and reptiles. P. 159-186. *In* C. L. Hubbs (ed.), *Zoogeography*. American Association for the Advancement of Science, Publ. 51.
- Peterson, E. A. 1966. Hearing in the lizard: Some comments on the auditory capacities of a nonmammalian ear. *Herpetologica* 22(3):161-171.
- Pope, C. H. 1955. The reptile world. A. A. Knopf Co., New York. xxvi + 325 + xiii p.
- Pough, F. H. 1969. Environmental adaptations in the blood of lizards. *Comp. Biochem. Physiol.* 31(6):885-901.
- Quay, W. B., and D. C. Wilhoft. 1964. Comparative and regional differences in serotonin content of reptilian brains. *J. Neurochem.* 11:805-811.
- Regal, P. J. 1966. Thermophilic response following feeding in certain reptiles. *Copeia* 1966(3):588-590.
- Retzlaff, E. W. 1949. The histology of the adrenal gland in the alligator lizard, *Gerrhonotus multicarinatus*. *Anat. Rec.* 105:19-28.
- Robinson, M. D., and T. R. Van Devender. 1973. Miocene lizards from Wyoming and Nebraska. *Copeia* 1973(4):698-704.
- Saint Girons, H. 1967. Morphologie comparée de la hypophyse chez les Squamata: Données complémentaires et rapport à la phylogénie des reptiles. *Ann. Sci. Natur. Zool.* (12)9:229-308.
- Saint Girons, M. 1970. Morphology of the circulating blood cells. P. 73-91. *In* C. Gans and T. S. Parsons (eds.), *Biology of the Reptilia*, Morphology C, vol. 3. Academic Press, New York.
- , and H. Saint Girons. 1969. Contribution à la morphologie comparée des érythrocytes chez les reptiles. *British J. Herpetol.* 4:67-82.
- Schmidt, K. P., and R. F. Inger. 1957. Living reptiles of the world. Doubleday and Co., New York. 287 p.
- Schultz, R., and M. Norberg. 1970. The effects of visual loss on lizard, *Gerrhonotus multicarinatus*, activity. *Psychon. Sci.* 19(1):10.
- Senn, D. G., and R. G. Northcutt. 1973. The forebrain and midbrain of some squamates and their bearing on the origin of snakes. *J. Morphol.* 140:135-152.
- Shaw, C. E. 1943. Hatching of the eggs of the San Diego alligator lizard. *Copeia* 1934(3):194.
- 1952. Notes on the eggs and young of some United States and Mexican lizards. *I. Herpetologica* 8(3):71-79.
- Skilton, A. J. 1849. Description of two reptiles from Oregon. *Amer. J. Sci. Arts* (2)7:202.
- Slater, J. R. 1963. Distribution of Washington reptiles. *Occas. Papers Dept. Biol. Univ. Puget Sound* (24):212-232.
- Smith, H. M. 1946. Handbook of lizards: Lizards of the United States and of Canada. Comstock Publ. Co., Ithaca, New York. xii + 557 p.
- , and E. H. Taylor. 1950. An annotated checklist and key to the reptiles of Mexico exclusive of the snakes. *U. S. Nat. Mus. Bull.* (199):1-253.
- Soule, J. D. 1970. The organic matrix in developing dental enamel and dentin studied by scanning electron microscopy. *Bull. Southern California Acad. Sci.* 69(2):104-111.
- Stebbins, R. C. 1944. Lizards killed by a millipede. *Amer. Midland Natur.* 32(3):777-778.
- 1948. Nasal structure in lizards with reference to olfaction and conditioning of the inspired air. *Amer. J. Anat.* 83(2):183-220.
- 1954. Amphibians and reptiles of western North America. McGraw-Hill Co., New York. xxiv + 528 p.
- 1958. A new alligator lizard from the Panamint Mountains, Inyo County, California. *Amer. Mus. Novitates* (1883):1-27.
- 1966. A field guide to western reptiles and amphibians. Houghton Mifflin Co., Boston. xiv + 279 p.
- Stejneger, L. 1893. Annotated list of the reptiles and batrachians collected by the Death Valley Expedition in 1891, with descriptions of new species. *N. Amer. Fauna* (7):159-228.
- Stephens, F. 1921. An annotated list of the amphibians and reptiles of San Diego County, California. *Trans. San Diego Soc. Natur. Hist.* 3(4):57-69.
- Storm, R. M. 1966. Amphibians and reptiles. Northwest. *Sci.* 40(4):138-141.
- Telford, S. R. 1965. A new nematode cestode from California lizards. *Japanese J. Exp. Med.* 35:301-303.
- Tihen, J. A. 1949. The genera of gerrhonotine lizards. *Amer. Midland Natur.* 41(3):580-601.
- Van Denburgh, J. 1897. The reptiles of the Pacific Coast and Great Basin. *Occas. Pap. California Acad. Sci.* (5):1-236.
- 1898. The *Gerrhonotus* of the San Lucan fauna, Lower California, with diagnosis of other west American species. *Proc. Acad. Natur. Sci. Philadelphia* 50:63-66.
- 1905. The reptiles and amphibians of the islands of the Pacific Coast of North America from the Farallons to Cape San Lucas and the Revilla Gigedos. *Proc. California Acad. Sci.* 3rd Ser. 4(1):1-40.
- 1922. The reptiles of western North America. Vol. 1. *Occas. Pap. California Acad. Sci.* 10:1-611.
- , and J. R. Slevin. 1914. Reptiles and amphibians of the islands of the west coast of North America. *Proc. California Acad. Sci.* 4th Ser. 4(5):129-152.
- , and — 1921. A list of the amphibians and reptiles of the peninsula of Lower California, with notes on the species in the collection of the Academy. *Ibid.* 4th Ser. 11(4):49-72.
- Waddick, J. W. 1963. A taxonomic study of gerrhonotine lizards. M. S. Thesis, University of Illinois, Urbana. 30 p.
- Wever, E. G. 1965. Structure and function of the lizard ear. *J. Auditory Res.* 5:331-337.
- Yarrow, H. C., and H. W. Henshaw. 1878. Report upon the reptiles and batrachians collected during the years 1875-1877 in California, Arizona and Nevada. *Ann. Rept. Sec. War for 1878, Part 3, Appendix L, in Appendix NN: 1628-1648.*

P. MIKE LAIS, MUSEUM OF NATURAL HISTORY, OREGON STATE UNIVERSITY, CORVALLIS, OREGON 97331.

Primary editor for this account, C. J. McCoy.

Published 5 November 1976 by the SOCIETY FOR THE STUDY OF OF AMPHIBIANS AND REPTILES.